

# Original WordList: TableName

## ID# Word

---

- 0 actinolite schist; emerald green, fine- to medium-grained, massive to well foliated; in part with up to 1 cm diameter weathered-out pits (after olivine ?); with locally abundant white feldspathic bands and veins; locally with anorthositic to gabbroic bands
- 1 amphibolite; fine- to medium-grained, heterogeneous, locally with light green calc-silicate streaks and pods; interpreted to be derived from a mafic volcanic protolith and/or mafic sills
- 2 calc-silicate gneiss; medium- to coarse-grained, heterogeneous, intercalated with grey garnetiferous psammite
- 3 diatexite derived from bp
- 4 diorite; medium grey, medium- to coarse-grained, massive to strongly lineated or foliated; locally grades into granodiorite. (u-pb zircon; 1858±3 ma)
- 5 felsic volcanogenic rocks; light grey to light greenish grey, very fine- to fine-grained, thinly laminated to massive, moderately foliated to mylonitic, locally interlayered with felsic to intermediate epiclastic volcanic rocks containing thin quartzitic
- 6 ferruginous psammite; pale reddish-brown to yellowish brown, fine- to medium- grained, well foliated, thin- to medium-bedded, rusty on weathered surfaces; locally with lean silicate facies iron formation and psammopelite intercalations
- 7 ferruginous psammite; pale reddish-brown, rusty on weathered surfaces, fine- to medium-grained, well foliated, heterogeneous; thin- to medium-bedded; locally interlayered with psammopelite, calc-silicate gneiss and cordierite-bearing felsic gneiss
- 8 granite; pink, medium- to coarse-grained, equigranular, massive to weakly foliated
- 9 granodiorite; light grey, medium- to coarse-grained, weakly to strongly foliated
- 10 hornblende ± actinolite ± cummingtonite schist; light grey, buff to rusty brown, fine- to medium-grained, thinly to thickly bedded, in part gossanous, in part graded bedding; intercalated with pebbly conglomerate derived mainly from volcanic detritus
- 11 intermediate paragneiss; tonalitic composition, migmatitic, well foliated, locally with clinopyroxene and orthopyroxene?cummingtonite bearing granitic leucosomes
- 12 intermediate volcanogenic rocks; medium grey to greenish grey, fine- to medium- grained, moderately foliated or mylonitic, massive to finely layered, hornblende porphyroblastic; possibly includes tuff, heterolithic tuff breccias, and polymictic conglomer
- 13 mafic volcanogenic rocks; greenish black to black, fine- to medium-grained, strongly foliated to mylonitic, homogeneous to banded, hornblende porphyroblastic; commonly interlayered with epidote- and carbonate-rich calc-silicate bands; locally includes ma
- 14 meta-arkose; pink, fine- to medium-grained, massive to moderately foliated, granoblastic; thinly to thickly bedded; locally with trough cross-bedding and conglomerate lag deposits
- 18 metagabbro; dark grey to greenish black, medium- to coarse-grained, inequigranular, mylonitic and commonly hornblende porphyroblastic
- 19 metapelite and metapsammite; grey, with biotite and garnet, fine- to medium- grained, migmatitic; intercalated with calc-silicate gneiss
- 15 meta-psammite; light grey to light pinkish grey, medium-grained, granoblastic, moderately to well foliated, thin to medium bedded; dominated by muscovite-bearing, white to pink pegmatite
- 16 meta-psammite; light grey to light pinkish grey, medium-grained, granoblastic, moderately to well foliated, thin to medium bedded; injected by 0 to 30 percent muscovite-bearing, white to pink pegmatite
- 20 metasedimentary and altered volcanogenic rocks
- 17 meta-turbidite; migmatitic, light grey, medium- to coarse-grained, with alternating pelitic and psammitic layers; locally with calc-silicate gneiss boudins
- 21 mixed diorite, granodiorite, tonalite, and granite; medium grey to light grey, fine- to medium-grained, equigranular, massive to strongly foliated and gneissic; locally migmatitic (u-pb zircon; 1891±4 ma tonalite and 1886±5 ma granite)
- 22 olivine websterite; massive, fine- to medium-grained
- 23 polymictic meta-conglomerate; light to dark grey, poorly sorted, matrix supported with strongly elongated pebble- to cobble-sized clasts of volcanic, plutonic, and sedimentary origin; matrix is fine- to medium-grained and biotite-hornblende to biotite-mu
- 24 psammopelite, pelite; medium grey, fine- to medium-grained, granoblastic, well foliated, poorly layered; locally includes minor calc-silicate gneiss, pale grey to light yellowish, laminated to massive, heterogeneous, fine- to medium-grained; pitted on we
- 25 psammopelitic schist and gneiss; light grey to buff, fine to medium-grained, granoblastic, well foliated, local graded bedding, typically migmatitic with 5-40 percent layer-parallel quartzitic to tonalitic leucosomes, locally cut by amphibolite dykes
- 26 pyroxenite (clinopyroxenite)?; dark green to brownish green, coarse-grained, massive to weakly foliated, equigranular
- 27 quartzofeldspathic paragneiss; beige to cream coloured, fine- to medium-grained, locally rusty
- 28 tonalite; medium grey to light grey, fine- to medium-grained, equigranular, massive to strongly foliated
- 29 ultramafic rock; emerald green, greenish black, to yellowish green, fine- to coarse-grained, massive to mylonitic, homogeneous to well banded